

## **PRESERVE** the option for organ donation!



- **DO NOT** terminally extubate before calling KODA
- **DO NOT** remove life-sustaining therapies before calling KODA (including pressor support, ECMO)
- **Allow KODA to initiate** Conversations about Organ and Tissue Donation

To maintain the option for **TISSUE** donation:

- Report all deaths to KODA **within 1 hour** of time of death
- **DO NOT** release the body to the funeral home

### PARAMETERS TO MAINTAIN:

- MAP between 60-110
- CVP 4-12
- EF  $\geq$  50%
- P:F Ratio  $\geq$  300
- Sodium  $\leq$  155
- Glucose  $\leq$  180
- Urine Output  $\geq$  .5 cc/kg/hr (over 4 previous hours)
- Low Dose Vasopressors 0-1

# Donation after Circulatory Death (DCD) Checklist

As you are *aware*, your patient's family has made the difficult decision to withdraw support.  
The OPO is working collaboratively with the family as DCD (Donation after Circulatory Death) has been authorized.  
Please refer to this checklist and mark the items off when completed.

## DONOR MANAGEMENT GOALS

Temp: 96.8–100.4 F (36–38 C)  
SBP: >100 mmHG <180 mmHG or  
MAP: >60  
HR: 60–140  
SpO2: >95%

**Document:** hourly vital signs, I&Os, medication administration

In collaboration with the attending physician/  
healthcare provider:

- ☐ Arterial Line Placement
- ☐ q6 Labs
- ☐ ABG as requested
- ☐ Other requested testing (CT, Echo, etc)

**4 HOURS PRIOR TO OR**